

## INDEX TO VOLUME IV

New names, and the final members of new combinations, are in **bold face type**

- Abies*, 91, 94, 145, 146, 184, 185, 190, 261; *balsamea*, 145, 149, 176, 177, 185, 187, 189, 192, 193; *canadensis*, 184; *concolor*, 145; *Fraseri*, 177; *grandis*, 94, 97, 145, 146; *lasiocarpa*, 58, 65, 145, 146; *magnifica*, 145; *nobilis*, 145, 146; *pectinata*, 177  
*Acer* *Negundo*, 266; *rubrum*, 328  
*Achlya*, 320-322, 326; *americana*, 319; 320; *DeBaryana*, 319, 320, 323, 324; *decorata*, 325; *glomerata*, 325, 326; *polyandra*, 319, 320, 323, 324; *prolifera*, 319, 320, 323, 324; *racemosa*, 325, 326; *racemosa spinosa*, 325; *racemosa stelligera*, 325; *spinosa*, 326  
*Achyla* *DeBaryana* Humphrey and the *prolifera* group, 319  
*Achlya glomerata* sp. nov., 325  
*Acolium*, 152  
*Actaea alba*, 9; *perennans*, 9  
*Adoxa*, 20; *Moschatellina*, 20, 32  
*Aecidium*, 13, 41, 195; *abundans*, 28; *auriellum*, 59; *elatinum*, 58; *monoiicum*, 59-61; *Polemonii*, 29; *sclerothecioides*, 20, 59  
*Aesculus glabra*, 11, 12, 52, 53  
*Agaricaceae* of the Pacific Coast, The —I, 205; II, 231; III, 294  
*Agaricaceae* of tropical North America, The—V, 72  
*Agarics*, New combinations for tropical, 331  
*Agaricum*, 273  
*Agaricus*, 239, 294; ***Abramsii***, 298; *alneus*, 213; *arenicola*, 81; *atrocoeruleus*, 214; ***bivelatoides***, 297; *bivelatus*, 296, 297; *bulbosus*, 295; *californicus*, 294; *campestris*, 239, 294, 296, 299; *campestris silvicola*, 295; *cerodes*, 81; ***comptuloides***, 297; *comptulus*, 298; *connatus*, 216; *copriniceps*, 81; ***crocodilinus***, 300; *diminutivus*, 298; *euthygrammus*, 76; *flammans*, 261; ***flavitingens***, 298; *haemorrhoidarius*, 295; *Herbarum*, 244; ***Hillii***, 298; ***hondensis***, 296; *limonellus*, 261; *liratus*, 248; ***McMurphyi***, 299; *multifidus*, 213; *oinodes*, 76; *popularis*, 80; *Pattersonae*, 296; *pectinatus*, 76; *pediades*, 76; *phalloides*, 240; *placomycetes*, 295; *porrigens*, 215; *radiatus*, 213; *semiorbicularis*, 76; *silvaticus*, 295; *silvicola*, 295; *solitarius*, 240; *subnitens*, 297; ***subrufescentoides***, 299  
*Agrimonia*, 192; *gryposepala*, 191  
*Agropyron repens*, 39; *tenerum*, 18, 31, 39  
*Agrostis alba*, 18, 31; *hyemalis*, 9  
*Alectoria*, 152; *loxensis*, 153; *loxensis atroalbicans*, 155  
*Aleuria umbrina*, 221  
*Aleuria*, Artificial cultures of *Ascobolus* and, 218  
*Alopecurus*, 180; *pratensis*, 179, 192  
*Alsine*, 145  
*Althaea rosea*, 18  
*Amanita*, 3, 231, 239; *bivolvata*, 241; *calytrata*, 241; *calyptroides*, 241; *calyptroderma*, 241; ***mexicana***, 332; *muscaria*, 239; *ocreata*, 240; ***pantherinoides***, 262; *phalloides*, 225; ***praegemmata***, 262; *rubescens*, 243; ***umbrinidisca***, 262; *virosa*, 242  
*Amanitopsis*, 3, 231, 239; *adnata*, 243; *velosa*, 239; *volvata*, 239  
*Ambrosia psilostachya*, 23; *trifida*, 11, 22  
*Amelanchier*, 25, 26, 56, 57, 62; *canadensis*, 63; *erecta*, 24-26, 32, 53, 56, 57, 62-64; *vulgaris*, 25, 32, 57, 61-65  
*American mycological literature*, Index to, 160, 227, 285, 333  
*Amoeba*, 322  
*Amorpha fruticosa*, 10, 12; *nana*, 10, 11, 13  
*Anaptychia*, 126  
*Andropogon*, 17, 199, 200; *furcatus*, 17, 31, 200, 204; *glomeratus*, 200; *scoparius*, 9, 17, 31, 200; *virginicus*, 17, 31, 200, 204  
*Anemone thalictroides*, 9  
*Anthoxanthum odoratum*, 11  
*Anthurus borealis*, 167  
*Apios Apios*, 10; *tuberosa*, 12  
*Aplanes*, 322  
*Apocynum cannabinum*, 9-13, 28  
*Aquilegia canadensis*, 9  
*Arabis*, 59-61, 65; *Drummondii*, 60; *Holboellii*, 10, 11, 20; *retrofracta*, 61  
*Aragallus Lamberti*, 23, 32  
*Arenaria lateriflora*, 186, 187, 193

# INDEX TO VOLUME IV

New names, and the final members of new combinations, are in **bold face type**

- Abies*, 91, 94, 145, 146, 184, 185, 190, 261; *balsamea*, 145, 149, 176, 177, 185, 187, 189, 192, 193; *canadensis*, 184; *concolor*, 145; *Fraseri*, 177; *grandis*, 94, 97, 145, 146; *lasiocarpa*, 58, 65, 145, 146; *magnifica*, 145; *nobilis*, 145, 146; *pectinata*, 177  
*Acer* *Negundo*, 266; *rubrum*, 328  
*Achlya*, 320-322, 326; *americana*, 319; 320; *DeBaryana*, 319, 320, 323, 324; *decorata*, 325; *glomerata*, 325, 326; *polyandra*, 319, 320, 323, 324; *prolifera*, 319, 320, 323, 324; *racemosa*, 325, 326; *racemosa spinosa*, 325; *racemosa stelligera*, 325; *spinosa*, 326  
*Achyla* *DeBaryana* Humphrey and the *prolifera* group, 319  
*Achlya glomerata* sp. nov., 325  
*Acolium*, 152  
*Actaea alba*, 9; *perennans*, 9  
*Adoxa*, 20; *Moschatellina*, 20, 32  
*Aecidium*, 13, 41, 195; *abundans*, 28; *auriellum*, 59; *elatinum*, 58; *monoiicum*, 59-61; *Polemonii*, 29; *sclerothecioides*, 20, 59  
*Aesculus glabra*, 11, 12, 52, 53  
*Agaricaceae* of the Pacific Coast, The —I, 205; II, 231; III, 294  
*Agaricaceae* of tropical North America, The—V, 72  
*Agarics*, New combinations for tropical, 331  
*Agaricum*, 273  
*Agaricus*, 239, 294; ***Abramsii***, 298; *alneus*, 213; *arenicola*, 81; *atrocoeruleus*, 214; ***bivelatoides***, 297; *bivelatus*, 296, 297; *bulbosus*, 295; *californicus*, 294; *campestris*, 239, 294, 296, 299; *campestris silvicola*, 295; *cerodes*, 81; ***comptuloides***, 297; *comptulus*, 298; *connatus*, 216; *copriniceps*, 81; ***crocodilinus***, 300; *diminutivus*, 298; *euthygrammus*, 76; *flammans*, 261; ***flavitingens***, 298; *haemorrhoidarius*, 295; *Herbarum*, 244; ***Hillii***, 298; ***hondensis***, 296; *limonellus*, 261; *liratus*, 248; ***McMurphyi***, 299; *multifidus*, 213; *oinodes*, 76; *popularis*, 80; *Pattersonae*, 296; *pectinatus*, 76; *pediades*, 76; *phalloides*, 240; *placomycetes*, 295; *porrigens*, 215; *radiatus*, 213; *semiorbicularis*, 76; *silvaticus*, 295; *silvicola*, 295; *solitarius*, 240; *subnitens*, 297; ***subrufescentoides***, 299  
*Agrimonia*, 192; *gryposepala*, 191  
*Agropyron repens*, 39; *tenerum*, 18, 31, 39  
*Agrostis alba*, 18, 31; *hyemalis*, 9  
*Alectoria*, 152; *loxensis*, 153; *loxensis atroalbicans*, 155  
*Aleuria umbrina*, 221  
*Aleuria*, Artificial cultures of *Ascobolus* and, 218  
*Alopecurus*, 180; *pratensis*, 179, 192  
*Alsine*, 145  
*Althaea rosea*, 18  
*Amanita*, 3, 231, 239; *bivolvata*, 241; *calyptata*, 241; *calyptatoides*, 241; *calyptroderma*, 241; ***mexicana***, 332; *muscaria*, 239; *ocreata*, 240; ***pantherinoides***, 262; *phalloides*, 225; ***praegemmata***, 262; *rubescens*, 243; ***umbrinidisca***, 262; *virosa*, 242  
*Amanitopsis*, 3, 231, 239; *adnata*, 243; *velosa*, 239; *volvata*, 239  
*Ambrosia psilostachya*, 23; *trifida*, 11, 22  
*Amelanchier*, 25, 26, 56, 57, 62; *canadensis*, 63; *erecta*, 24-26, 32, 53, 56, 57, 62-64; *vulgaris*, 25, 32, 57, 61-65  
*American mycological literature*, Index to, 160, 227, 285, 333  
*Amoeba*, 322  
*Amorpha fruticosa*, 10, 12; *nana*, 10, 11, 13  
*Anaptychia*, 126  
*Andropogon*, 17, 199, 200; *furcatus*, 17, 31, 200, 204; *glomeratus*, 200; *scoparius*, 9, 17, 31, 200; *virginicus*, 17, 31, 200, 204  
*Anemone thalictroides*, 9  
*Anthoxanthum odoratum*, 11  
*Anthurus borealis*, 167  
*Apios Apios*, 10; *tuberosa*, 12  
*Aplanes*, 322  
*Apocynum cannabinum*, 9-13, 28  
*Aquilegia canadensis*, 9  
*Arabis*, 59-61, 65; *Drummondii*, 60; *Holboellii*, 10, 11, 20; *retrofracta*, 61  
*Aragallus Lamberti*, 23, 32  
*Arenaria lateriflora*, 186, 187, 193

- Armillaria, 207, 211, 238; *albolanaripes*, 211; *arenicola*, 212; *mag-nivellaris*, 212; *melles*, 211, 266; *nardosmia*, 238; *subannulata*, 211  
 Arnica, 10, 11, 13, 20, 22, 23, 27  
 Aronia, 25; *arbutifolia*, 25, 32, 63, 65; *nigra*, 25, 32  
 Artemisia, 21, 32; *dracunculoides*, 16, 21, 31, 32; *ludoviciana*, 21  
 Arthur, J. C., Cultures of Uredineae in 1910, 7; in 1911, 49  
 Artificial cultures of *Ascobolus* and *Aleuria*, 218  
*Asclepias syriaca*, 19, 31  
*Ascobolus*, 218, 220, 221; *applanatus*, 219; *carbonarius*, 221; *laeviporus*, 219; *latus*, 219; *magnificus*, 218, 219, 221, 222; *major*, 219; *miniatus*, 45, 47; *sarawacensis*, 219  
*Ascobolus* and *Aleuria*, Artificial cultures of, 218  
*Aspergillus*, 279; *flavescens*, 279, 280; *flavus*, 279; *fumigatus*, 279; *glauco-*  
*niger*, 279  
*Aspergillus* infecting *Malacosoma* at high temperatures, 279  
*Aster*, 7, 16, 21, 22, 144, 181, 199; *acuminatus*, 180, 181, 193; *adscen-*  
*dens*, 16, 30; *Drummondii*, 9, 21; *ericoides*, 19, 21, 31, 32; *multiflorus*, 19, 31; *Novae-Angliae*, 19, 31; *paniculatus*, 10, 13, 15, 21, 28, 32, 33; *punicus*, 181, 192; *Tweedyi*, 16  
*Astragalus*, 24; *canadensis*, 13; *carolinianus*, 23, 32; *sulphureus*, 23, 32  
*Atestia loxensis*, 153  
*Atheropogon curtispindulus*, 19, 31  
*Atriplex*, 54, 55, 198; *confertifolia*, 52; *hastata*, 54, 55, 64; *patula*, 179, 198  
*Aurantiporellus alboluteus*, 91  
*Auricularia auricula*, 40  
*Auriscalpium Auriscalpium*, 330  
*Bacidia*, 126; *endoleuca*, 131; *rubella*, 131; *subgranulosa*, 131  
*Bacterium Savastanoi*, 104  
*Baeomyces absolutus*, 132; *erythrel-*  
*lus*, 132  
*Bagniesiella*, 70; *australis*, 70; *Di-*  
*antherae*, 70  
 Banker, H. J., Type studies in the *Hydnaceae*—I. The genus *Manina*, 271; II. The genus *Steccherinum*, 309  
 Barlaea, 45, 47  
 Barlaeina, 45, 47  
*Berberis vulgaris*, 11, 18, 31  
 Beta, 198  
*Biatora*, 126, 130-132; *amorpho-*  
*carpa*, 127; *atrogrisea*, 131; *aurigera*, 127; *coarctata*, 127; *endocaeerulea*, 128; *lanuginosa*, 128; *oncodes*, 128; *subgranulosa*, 131; *triseptata artytoides*, 130  
*Bilimbia*, 126; *artytoides*, 130, 131; *hypnophila*, 130; *molybdis*, 130; *pallidissima*, 130; *radicicola*, 130; *sphaeroides*, 130; *terrestris*, 130; *thysanota*, 130  
*Bjerkandra adusta*, 91  
 Black knot disease of *Dianthera americana* L., A, 66  
*Boehmeria cylindrica*, 10, 17, 28  
*Bolbitis*, 73; *fragilis*, 73; *jalapensis*, 332; *mexicanus*, 332; *radians*, 246; *villipes*, 73  
 Boletaceae of the Pacific Coast, *Poly-*  
*poraceae* and, 91  
*Boletellus Ananas*, 99  
*Boletus Lakei*, 97; *luteus*, 98; *mira-*  
*bilis*, 217; *oregonensis*, 217; *Zelleri*, 217  
 Bolls, *Botryosphaeria* on cotton, 34  
*Boltonia asteroides*, 10, 12, 20  
*Bombyliospora tuberculosa*, 131  
*Botryosphaeria*, 34, 36; *Berengeriana*, 34; *fuliginosa*, 34, 36  
*Botryosphaeria* on cotton bolls, 34  
*Bryophagus Gloeocapsa*, 127  
*Buellia stipitata*, 139; *subdiscifor-*  
*mis*, 139  
 Bursa Bursa-pastoris, 55  
*Caecoma*, 30, 146, 187-189, 195; *Abietis-*  
*canadensis*, 188, 189, 193; *coni-*  
*geneum*, 146; *occidentale*, 30  
*Caecomurus*, 195  
*Calamagrostis canadensis*, 18, 31  
*Calamovilfa longifolia*, 18, 31  
 Calcium, 152  
*Calliospora*, 282  
*Callirrhoe involuerata*, 18, 31, 53  
*Calocera cornea*, 40  
*Calvatia gigantea*, 330  
*Calyptospora*, 146; *columnaris*, 146, 177, 192  
*Capsella*, 198  
*Carduus*, 23; *Flodmanii*, 11, 22, 23, 32  
*Carex*, 14-16, 21, 28, 54, 180, 199, 201; *arctata*, 9, 180, 193; *aristata*, 17, 31; *brunnescens*, 28; *crinita*, 180, 193; *debilis* Rudgei, 180, 193; *deflexa*, 21, 22, 32, 181, 192; *festiva*, 16, 30; *Goodenovii*, 28, 33; *gracillima*, 14, 201, 204; *intumescens*, 21, 22, 32, 180, 181, 192, 193, 199, 204; *lanuginosa*, 15, 30, 54, 64; *pallidescens*, 13, 14, 30; *Pennsylvanica*, 7, 9; *pubes-*  
*cens*, 14, 200, 201, 204; *scoparia*, 15, 22, 30, 181, 182, 192, 193, 199; *siccata*, 16, 31; *stellulata*, 9; *steno-*

- phylla, 16, 31; stipata, 181, 193; stricta, 17, 31; tenella, 9; tenuis, 13, 14, 30, 180, 201; tribuloides, 199, 204; trichocarpa, 15, 30; trisperma, 9, 181, 193; varia, 22  
 Cassia chamaecrista, 11, 12  
 Castanea dentata, 170, 171  
 Castilleja, 142; integra, 142  
 Catalpa, *Polystictus versicolor* as a wound parasite of, 263  
 Catillaria **leptochella**, 128; **rosea**, 128  
 Caulophyllum thalictroides, 9  
 Cephalanthus occidentalis, 19, 31  
 Cerastium oreophilum, 58, 65  
 Cerastium, 145  
 Ceratitium, 195  
 Ceromyces communis, 98, 100; **mirabilis**, 98, 217; **oregonensis**, 99, 217; viscidus, 99; **Zelleri**, 99, 217  
 Certain species of Puccinia and Uromyces, Correlation between, 194  
 Chamaedaphne calyculata, 178, 192  
 Chanterel, 206; alektorolophoides, 206; behringensis, 206; bryophilus, 206; Chantarellus, 207; floccosus, 206; infundibuliformis, 206  
 Cheiranthus Menziesii, 59  
 Chelone glabra, 9  
 Chenopodium, 198; album, 18, 19, 31, 54, 55, 64, 179, 192, 198  
 Chestnut and oak, The large leaf spot of, 170  
 Chrysopogon avenaceus, 10  
 Cicuta maculata, 178, 179, 192  
 Cladonia, 132; aggregata, 132; alpestris, 132; angustata, 132; ceratophylla, 132; dactylota, 132; degenerans, 132; didyma muscigena, 132; fimbriata simplex, 132; fimbriata subulata, 132; Floerkeana intermedia, 132; pityrea, 132; rangiformis, 132; sylvatica, 132  
 Clavaria, 271, 273, 275  
 Clematis virginiana, 11  
 Cleome, 54, 198; spinosa, 54, 55, 64  
 Clitocybe, 4, 330  
 Clitopilus abortivus, 6; **Earlei**, 332  
 Coast, Polyporaceae and Boletaceae of the Pacific, 91  
 Coast, The Agaricaceae of the Pacific—I, 205; II, 231; III, 294  
 Coccocarpia pellita, 135; pellita cro-nia, 135  
 Coelocaulon divergens, 153  
 Coenogonium Leprieurii, 127  
 Coker, W. C., Achlya DeBaryana Humphrey and the proliferata group, 319; Achlya glomerata sp. nov., 325; & Hyman, O. W., Thraustotheca clavata, 87  
 Coleosporium, 144; Vernoniae, 29, 33, 57, 65  
 Collybia, 4; stipitaria, 5  
 Collybidium dryophilum, 164; zonatum, 4  
 Coltricia perennis, 92  
 Comandra umbellata, 17, 31  
 Combinations for tropical agarics, New, 331  
 Coniophora cerebella, 223  
 Conocybe, 72, 74, 244, 247; **angusticeps**, 248; **Bryorum**, 247, 248; **echinospora**, 75, 332; **Hypnorum**, 75, 247; **lirata**, 248; **semilanceata**, 248; **Sphagnorum**, 248; tener, 74, 247  
 Convallaria majalis, 11  
 Coprinus, 223; Brassicae, 2; Spraguei, 5  
 Cora pavonia, 140  
 Coriellus Sepium, 92, 268, 270  
 Coriolus abietinus, 92; nigromarginatus, 92; prolificans, 92; versicolor, 92; **washingtonensis**, 92, 217  
 Cornicularia loxensis, 153  
 Correlation between certain species of Puccinia and Uromyces, 194  
 Cortinari, 72, 81, 243; cinnamomeus, 81; **mexicanus**, 81; Sitenisii, 81  
 Corydalis aurea, 18; montanum, 55; sempervirens, 19  
 Cotton bolls, Botryosphaeria on, 34  
 Coursetia glandulosa, 282, 283  
 Crataegus, 25, 261; cernonis, 26, 32, 57, 62, 65; punctata, 25, 32; tomentosa, 24, 25, 32, 57, 63, 65  
 Craterellus cornucopioides, 5  
 Crepidopus, 207, 216; **connatus**, 216; ostreatus, 216; **serotinus**, 216; **subsapidus**, 216, 217  
 Crepidotus, 244; calolepis, 245; fulvotomentosus, 245; Herbarum, 244; malachias, 245; mollis, 245; puberulus, 245; **submollis**, 245  
 Cronartium, 144; coleosporioides, 142; Quercum, 144; Quercus, 26, 33; ribicola, 42, 104, 143  
 Crouania, 45, 47; miniata, 45  
 Cryptoporus volvatus, 96  
 Cultures of Ascobolus and Aleuria, Artificial, 218  
 Cultures of heteroecious rusts, 175  
 Cultures of Uredineae in 1910, 7; in 1911, 49  
 Cummings, Clara Eaton, in Jamaica, An enumeration of lichens collected by—I, 125  
 Cydonia vulgaris, 25, 26, 32, 56  
 Dacryomyces, 40; deliquescent, 40  
 Daedalea unicolor, 328

- Delphinium formosum, 10; scaposum, 11; tricornis, 11
- Descriptions of two new species, The genus Lamprospora, with, 45
- Dianthera, 66; americana, 66, 70
- Dianthera americana L., A black knot disease of, 66
- Diaporthe parasitica, 157
- Dichodium byrsinum, 133
- Dichroum, 311
- Dictyolus, 206; retirugus, 206
- Dictyophora duplicata, 167
- Dictyuchus, 87, 322; clavatus, 87
- Diplodia, 34, 36; gossypina, 34, 35
- Diploschistes scruposus, 127
- Dirca palustris, 9-12
- Disease of Dianthera americana L., A black knot, 66
- Diseases in the eastern states, Notes upon tree, 148
- Distichlis, 179; maritima, 202; spicata, 12, 18, 31, 52, 54, 55, 64, 179, 192, 198, 204
- Distribution, Oropogon loxensis and its North American, 152
- Dodge, B. O., Artificial cultures of Ascobolus and Aleuria, 218
- Doellingeria, 16; umbellata, 15
- Drosophila, 294, 302; **ambigua**, 304; appendiculata, 302; **atrofolia**, 303; **californica**, 304; **campanulata**, 304; **longipes**, 303
- Dryodon, 272, 276
- Eastern states, Notes upon tree diseases in the, 148
- Edgerton, C. W., Botryosphaeria on cotton bolls, 34
- Elaeagnus angustifolia, 11, 13
- Eleocharis, 203; palustris, 12
- Elephantopus carolinianus, 29
- Elfvingia fasciata, 105; megaloma, 96
- Elymus canadensis, 11; triticoides, 39
- Enumeration of lichens collected by Clara Eaton Cummings in Jamaica, An—I, 125
- Epilobium, 177; angustifolium, 176, 192
- Erigeron, 7, 13, 16, 22, 199; annuus, 13
- Erioderma, 135; **microcarpa**, 134; Wrightii, 134
- Eupatorium perfoliatum, 203; serotinum, 12
- Euphorbia, 202, 203; commutata, 13; corollata, 202; Cyparissias, 23, 24
- Eurodium repens, 279
- Euthamia, 16, 22; graminifolia, 15, 30
- Evolvulus pilosus, 28, 33
- Favolus, 317
- Festuca confinis, 27, 33
- Flammula, 250; astragalina, 251; californica, 253; carbonaria, 165, 256; **decorata**, 262; **echinulispora**, 262; fulvella, 253; **Hillii**, 262; **laeticolor**, 262; **lata**, 262; **ornatula**, 262; **pallida**, 262; penetrans, 254; **permollis**, 262; sapinea, 254; **spinulifer**, 262; **spumosa**, 254; **subcarbonaria**, 262; **subflavida**, 262; **vialis**, 262; **viridans**, 262; **viscidissima**, 262
- Fomes, 109; annosus, 96, 105; Earlei, 109-111, 114; fraxinophilus, 105; juniperinus, 109, 110, 114; Laricis, 95, 97, 105; roseus, 97; **texanus**, 109, 110, 112, 114; unguis, 97
- Forest trees, Notes on some western Uredineae which attack, 141
- Fraser, W. P., Cultures of heterocyclic rusts, 175
- Fraxinus lanceolatus, 11
- Friesites, 272, 276
- Fungi, Illustrations of—X, 1; XI, 163; XII, 289
- Fusarium, 104, 106; oxysporium, 104
- Galera, 74, 247; angusticeps, 248; Besseyi, 248; Bryorum, 247; cubensis, 75; **echinospora**, 332; grisea, 75; Hypnorum, 75; lirata, 249; semilanceata, 248; simulans, 74; sphaerobasis, 247; Sphagnorum, 248; tener, 74; versicolor, 247
- Gaura biennis, 15
- Geaster **juniperinus**, 85, 86; marginatus, 84-86; minimus, 84-86
- Geaster minimus Schw. and its relatives. Notes on Iowa saprophytes—I, 84
- Gee, W. P., & Massey, A. B., Aspergillus infecting Malacosoma at high temperatures, 279
- Gentiana, 203
- Genus Lamprospora, with descriptions of two new species, The, 45
- Genus Lasiosphaeria, The, 115
- Genus Manina, The. Type studies in the Hydnaceae—I, 271
- Genus Steccherinum, The. Type studies in the Hydnaceae—II, 309
- Geopetalum, 207, 214; **densifolium**, 215, 217; geogenium, 214; **oregonense**, 215, 217; **porrigens**, 215; **subsepticum**, 215, 217
- Glaux maritima, 187
- Gloeophyllum hirsutum, 97
- Godronia, 123; Bethell, 123; striata, 123
- Gomphidius, 260, 294, 306, 308; nigricans, 306; oregonensis, 306; **tomentosus**, 307; vinicolor, 307, 308

- Graves, A. H., The large leaf spot of chestnut and oak, 170
- Grindelia squarrosa*, 10, 19, 20, 22, 27, 31
- Group, *Achlya* DeBaryana Humphrey and the *prolifera*, 319
- Gyalecta*, 127; *Gloeocapsa*, 127
- Gymnoconia interstitialis*, 53
- Gymnopilus*, 244, 250, 261; **californicus**, 252; **carbonarius**, 250, 256, 257; **decoratus**, 250-252, 262; **echinulisporus**, 250, 255, 262; **foedatus**, 257; **fulvellus**, 253; **Hillii**, 253, 262; **laeticolor**, 250, 251, 262; **latus**, 257, 262; **ornatus**, 251, 262; **pallidus**, 252, 262; **penetrans**, 254; **permollis**, 252, 262; **sapineus**, 254; **spinulifer**, 254, 262; **spumosus**, 254, 256; **subcarbonarius**, 256, 262; **subflavidus**, 250, 252, 262; **vialis**, 255, 262; **viridans**, 250, 257, 262; **viscidissimus**, 256, 262
- Gymnosporangium*, 24, 40, 41, 50, 62, 63, 106, 195, 197, 223; *bermudianum*, 50; *Betheli*, 25, 32; *clavariaeforme*, 24, 32, 53, 56, 64; *clavipes*, 24, 32; *cornutum*, 25, 32; *Davisii*, 25, 32; *durum*, 61, 62, 65; *effusum*, 50, 62, 65; *Ellisii*, 50; *fraternum*, 50; *gracile*, 63, 65; *inconspicuum*, 57, 64; *juvenescens*, 26, 62; *juniperinum*, 57, 65; *Juniperi-virginianae*, 24, 32; *Kernianum*, 62, 65; *Libocedri*, 57, 65; *Nelsoni*, 26, 32, 61, 62, 65; *nidus-avis*, 25, 26, 32, 56, 64; *speciosum*, 63; *tremelloides*, 57, 65
- Haematomma puniceum*, 138
- Halameda*, 320
- Hebeloma*, 72, 82, 243; **Broadwayi**, 82; **cinchonense**, 82; *foedatum*, 257; *longicaudum*, 83; **subincarnatum**, 83
- Hedgcock, G. G., Notes on some western Uredineae which attack forest trees, 141; & Long, W. H., Preliminary notes on three rots of juniper, 109
- Hericium*, 272-275; *alpestre*, 274, 275; *Echinus*, 274; *grandis*, 277; *Hystrix*, 274; *Notarisii*, 274
- Heteroecious* rusts, Cultures of, 175
- Heterothecium*, 128, 129, 131
- Hevea*, 102
- Hibiscus militaris*, 18
- Hicoria*, 121
- High temperatures, *Aspergillus* infecting *Malacosoma* at, 279
- Hordeum jubatum*, 39
- Howe, R. H., Jr., *Oropogon loxensis* and its North American distribution, 152
- Humaria calospora*, 48
- Hydnaceae, Type studies in the—I. The genus *Manina*, 271; II. The genus *Steccherinum*, 309
- Hydnum*, 274, 276, 317; *abietinum*, 276; *alpestre*, 275; *basi-asperatum*, 312, 313; *Caput-medusae*, 274, 275, 277; *Caput-ursi*, 272, 277; *conchiforme*, 318; *coralloides*, 272-276; *croceum*, 278; *Daviesii*, 309; *decurrens*, 318; *dichroum*, 310-312, 330; *Echinus*, 274; *Erinaceus*, 272, 275, 277, 330; *flabelliforme*, 311, 312; *glabrescens*, 315; *glabrescens*, 312, 315, 316; *guaraniticum*, 312, 313; *hystricinum*, 273, 274; *Hystrix*, 273, 274; *laciniatum*, 276; *laeticolor*, 316; *Notarisii*, 275; *ochraceum*, 309, 310, 312; *Persoonianum*, 312; *plumarium*, 317, 318; *plumosum*, 317; *pudorium*, 310-312; *pusillum*, 313; *ramarium*, 274; *ramosum*, 276; *rawakense*, 312, 313, 315; *reniforme*, 312; *Rhois*, 311, 312; *ruginosum*, *Necium Farlowii*, 182, 193
- Hydrocybe*, 207; *albo-umbonata*, 332; *arenicola*, 208, 217; *aurantia*, 332; *coccinea*, 208; *conica*, 207; **constans**, 208, 217; **cremicolor**, 209, 217; *Earlei*, 332; *flavolutea*, 332; *hondurensis*, 332; *rosea*, 332; *subcaespitosa*, 332; *subflavida*, 332; *subminiata*, 332; *troyana*, 332
- Hydrophyllum capitatum*, 10, 11, 27, 29, 53; *virginicum*, 10, 12, 20
- Hygrophorus*, 207, 209; **albo-umbonatus**, 332; **arenicola**, 217; **aurantius**, 332; **coccineus**, 208; **constans**, 217; **cremicolor**, 217; *Earlei*, 332; *eburneus*, 209; *flavodiscus*, 6; **flavoluteus**, 332; **fragrans**, 210; *hondurensis*, 332; *Laurae*, 210; *limacinus*, 210; *miniatus*, 208; **roseus**, 332; *rubropunctatus*, 210; **subcaespitosus**, 332; **subflavidus**, 332; **subminiatus**, 332; **subpustulatus**, 210; **troyanus**, 332; **varicolor**, 209
- Hyman, O. W., Coker, W. C. & *Thraustotheca clavata*, 87
- Hypholoma*, 294, 302, 303, 305; *ambiguum*, 304; *appendiculatum*, 1, 302; *atrofolium*, 303; *californicum*, 304; *campanulatum*, 304; *capnoides*, 305; *cutifractum*, 302; *fasciculare*, 305, 306; *flocculentum*, 303; *lacrymabundum*, 302; *longipes*, 303, 304; *sublateritium*, 302
- Hypodendrum*, 72, 244, 261; **flam-**

- mans**, 261; **limonellum**, 261; **oregonense**, 261, 262  
*Hypodermium*, 101, 195  
*Hyponovris*, 207, 213; *alneus*, 213, 214  
*Hypoxylon*, 121; *miaceum*, 121
- Illustrations of fungi—X, 1; XI, 163; XII, 289  
*Impatiens*, 20; *aurea*, 11, 20, 32  
 Index to American mycological literature, 160, 227, 285, 333  
*Inocybe*, 72, 82, 243; **jamaicensis**, 82; *rimosa*, 4  
*Inonotus dryophilus*, 104; *texanus*, 104  
 Iowa saprophytes, Notes on—I. Geaster minimus Schw. and its relatives, 84  
*Iris*, 28; *versicolor*, 28  
*Irpiciporus*, 94  
*Ischnoderma fuliginosum*, 92  
*Isopyrum bitermum*, 9, 11
- Jamaica, An enumeration of lichens collected by Clara Eaton Cummings in—I, 125  
*Juncus balticus*, 22, 23, 32  
 Juniper, Preliminary notes on three rots of, 109  
*Juniperus monosperma*, 63, 65, 112, 113; *sabinoides*, 112, 113; *scopulorum*, 25, 32; *sibirica*, 24, 25, 32, 56, 57, 64, 65; *utahensis*, 56, 61, 62, 64, 65, 112, 113; *virginiana*, 24–26, 32, 56, 62, 64, 65, 85, 109, 111
- Kalmia*, 150; *angustifolia*, 150; *latifolia*, 150  
 Knot disease of *Dianthera americana* L., A black, 66  
*Koeleria*, 59, 60; *cristata*, 20, 31, 59
- Laccaria*, 207, 211; *laccata*, 211; *ochropurpurea*, 211  
*Lachnea scutellata*, 38  
*Laciniaria punctata*, 9, 10, 57; *scariosa*, 12, 20, 29, 57; *spicata*, 9, 10, 12  
*Lactaria*, 207, 289  
*Lactuca canadensis*, 16, 31; *sativa*, 16, 31; *scariola*, 13  
*Laetiporus speciosus*, 92  
*Lamprospora*, 45, 47; **areolata**, 48; **tuberculata**, 47  
*Lamprospora*, with descriptions of two new species, The genus, 45  
 Large leaf spot of chestnut and oak, The, 170  
*Larix*, 29, 188, 189; *laricina*, 29, 58, 188, 189  
*Lasiosphaeria*, 115–117, 123; *ambigua*, 122; *canescens*, 118; *dichrospora*, 117, 123, 124; **globularis**, 117, 121, 123; *hirsuta*, 119; *hispida*, 117, 119–121, 123; *Hystrix*, 118; **jamaicensis**, 117, 122, 123; **mucida**, 117, 118, 121, 124; **multiseptata**, 117, 120, 124; *newfieldiana*, 117, 122, 123; *ovina*, 118; *Rhacodium*, 119; *spermoides*, 121; *striata*, 123; *strigosa*, 117, 118, 123; *stuppea*, 117, 119, 123; *terrestris*, 117, 120, 124  
*Lasiosphaeria*, The genus, 115  
*Lathyrus palustris*, 13  
 Leaf spot of chestnut and oak, The large, 170  
*Lecanora*, 138; *pallida*, 137; *subfusca*, 137; *varia*, 138  
*Lecidea*, 126–131; *leucoblephara*, 126; *microphyllina subgranulosa*, 131  
*Ledum*, 26, 178; *groenlandicum*, 26, 33, 177, 178, 192  
*Lentinus lepideus*, 257  
*Lenzites betulina*, 97; *sepiaria*, 101  
*Lepargyrea canadensis*, 10–13  
*Lepidium*, 198; *virginicum*, 55  
*Lepiota*, 231, 238; *adnatifolia*, 237; **agricola**, 332; *amianthina*, 237, 238; **amplifolia**, 233; *aspera*, 238; *asperula*, 238; *brunnescens*, 236; *carcharias*, 237; **castaneidiscas**, 232; **castanescens**, 234; *cepaestipes*, 232; **concentrica**, 235; *cretaea*, 232; *cristata*, 232, 233; *cristatella*, 232; *eriphora*, 238; **fuliginescens**, 236; *fulvodisca*, 212; **fumosifolia**, 233; *fuscusquamia*, 235; *granosa*, 237; *granulosa*, 237; *granulosa amianthina*, 238; **mag-nispora**, 237; **McMurphyi**, 217; *mutata*, 233; *nardosmioides*, 238; *naucina*, 236; *petasiformis*, 232; **roseicremea**, 217; **roseifolia**, 235; **roseilivida**, 234; *rubrotincta*, 237; **rubrotinctoides**, 236; *rugoso-reticulata*, 237; **Sequoiarum**, 233; **subfelina**, 234; **subnivosas**, 231  
*Leptogium bullatum*, 133; *chlorome-lum*, 133; *foveolatum*, 134; *lacerum*, 133; *marginellum*, 133; *myochroum tomentosum*, 134; *phyllocarpum*, 133; *punctulatum*, 133; *saturninum*, 134; *tremelloides*, 134; *tremelloides impresso-punctatum*, 134  
*Leptonia atosquamosa*, 332; **cin-chonensis**, 332; **Earlei**, 332; **mexicana**, 332  
*Leptoniella atosquamosa*, 332; *cin-chonensis*, 332; **Earlei**, 332; *mexicana*, 332  
*Leptospora*, 115, 117; *ovina*, 118; *spermoides*, 121; *strigosa*, 118  
*Leucomyces*, 239; *mexicanus*, 332  
 Lewis, I. M., A black knot disease of *Dianthera americana* L., 66



- Libocedrus decurrens*, 57, 65  
 Lichens collected by Clara Eaton Cummings in Jamaica, An enumeration of—I, 125  
*Limacella*, 207, 212; *agricola*, 332; *fulvodisca*, 212; **McMurphyi**, 213, 217; *roseicremea*, 212, 217  
 Literature, Index to American mycological, 160, 227, 285, 333  
*Lithospermum canescens*, 10, 11  
*Lobaria*, 135; *corrosa*, 135; *erosa*, 136; *pallida*, 135; *peltigera*, 135; *quercizans*, 136  
*Locellina*, 244, 262; *californica*, 262; *stercoraria*, 262  
 Long, W. H., Two new species of rusts, 282  
 Long, W. H., Hedgecock, G. G. & Preliminary notes on three rings of juniper, 109  
*Lopadium amaaurum*, 131; *leucoxanthum*, 131  
*Lophodermium nervisequum*, 149  
*Lycopus americanus*, 17, 31, 54, 64  
*Lygodesmia juncea*, 58, 65  
*Lysimachia quadrifolia*, 9, 10; *terrestris*, 9, 12, 13, 22, 28  
 Macbride, T. H., Notes on Iowa saprophytes—I. *Geaster minimus* Schw. and its relatives, 84  
*Macrocalyx Nyctelea*, 11, 28  
*Macrophoma*, 34, 103  
*Mahonia Aquifolium*, 11  
*Malacosoma americana*, 279, 280  
*Malacosoma* at high temperatures, *Aspergillus* infecting, 279  
*Malus*, 56; *coronaria*, 26, 56, 63, 64; *Malus*, 24, 32, 63  
*Malvastrum coccineum*, 52  
*Manina*, 271-276, 278; **Caput-ursi**, 277; *coralloides*, 276, 330; *cordiformis*, 272, 275, 277, 330; *flagellum*, 276; **Schieder-mayeri**, 277  
*Manina*, The genus. Type studies in the Hydnaceae—I, 271  
*Marasmius*, 4, 5, 157; *magnisporus*, 166; *oreades*, 224; *Vaillantii*, 167; *viticola*, 167  
*Martela*, 273  
*Martella*, 273-275; *Echinus*, 273, 274  
 Massey, A. B., Gee, W. P., & *Aspergillus* infecting *Malacosoma* at high temperatures, 279  
*Medicago*, 24, 56; *sativa*, 13, 24, 32, 56, 64  
*Medusina*, 272, 275; *coralloides*, 276; *patula*, 272, 275, 277  
*Megalospora Cummingsiae*, 129; *jamaicensis*, 129; *sulphurata*, 129; *versicolor*, 129  
*Melampsora*, 30, 146, 188, 189; *albertensis*, 29, 33, 58, 65; *arctica*, 187, 193; *Bigelowii*, 189; *Medusae*, 188, 189, 193  
*Melampsorella elatina*, 58, 65, 145  
*Melampsoropsis*, 175; *abietina*, 26, 33, 178, 192; *Cassandrae*, 178, 192; *ledicola*, 177, 192; *Pyrolae*, 183, 184, 193  
*Melanoleuca jalapensis*, 332; *jamaicensis*, 332; *subisabellina*, 332  
*Melanops Quercuum*, 34  
*Meriolix*, 15; *serrulata*, 9, 15, 30, 54  
*Merisma*, 274  
*Merulius lacrymans*, 223, 327  
*Metasphaeria sublanosa*, 118  
*Microsphaera quercina*, 223  
*Microphiale lutea*, 127  
*Mimulus ringens*, 10, 12  
*Monochaetia*, 172; *Desmazierii*, 104, 170-174, 330, 331; *monochaeta*, 172; *pachyspora*, 170, 331  
*Monolepis Nuttalliana*, 18, 19, 54  
*Mucor*, 88  
*Muhlenbergia gracilis*, 18; *racemosa*, 18, 31  
 Murrill, W. A., Illustrations of fungi—X, 1; XI, 163; XII, 289; New combinations for tropical agarics, 331; *Polyporaceae* and *Boletaceae* of the Pacific Coast, 91; The *Agaricaceae* of the Pacific Coast—I, 205; II, 231; III, 294; The *Agaricaceae* of tropical North America—V, 72  
*Mycelophagus Castaneae*, 223  
*Mycena*, 72, 73; *fragilis*, 73; *jalapensis*, 73, 332; *mexicana*, 73, 332; *praedecurrens*, 165; *vexans*, 168; *villipes*, 73; *vulgaris*, 165  
 Mycological literature, Index to American, 160, 227, 285, 333  
*Myrica cerifera*, 9, 10, 11, 28  
*Myxosporium acerinum*, 151  
*Naematelia encephala*, 40; *nucleata*, 40  
*Napaea dioica*, 18  
*Naucoria*, 72, 74, 75, 81, 243; *corticola*, 77; *cyathicola*, 77; *Earlei*, 77; *euthugrammus*, 76; *hepaticicola*, 78; *jalapensis*, 77; *montana*, 78; *oinodes*, 76; *paludosa*, 249; *pectinata*, 76; *pediades*, 76; *pellucida*, 78; *Sacchari*, 79; *semi-orbicularis*, 76; *sideroides*, 80; *spinulifer*, 79; *subvelosa*, 164; *tepeitensis*, 79; *Underwoodii*, 80; *xuchilensis*, 80  
*Necium Farlowii*, 182, 193  
*Nectria cinnabarina*, 266  
 New combinations for tropical agarics, 331



- New species of rusts, Two, 282  
 New species, The genus *Lamprospora*, with descriptions of two, 45  
 News and notes, 37, 101, 157, 223, 327  
 Nigredo, 195, 223  
 North America, The Agaricaceae of tropical—V, 72  
 North American distribution, *Oropogon loxensis* and its, 152  
 Notes, News and, 37, 101, 157, 223, 327  
 Notes on Iowa saprophytes—I. *Geaster minimus* Schw. and its relatives, 84  
 Notes on some western Uredineae which attack forest trees, 141  
 Notes on three rots of juniper, Preliminary, 109  
 Notes upon tree diseases in the eastern states, 148  
 Oak, The large leaf spot of chestnut and, 170  
*Ocellularia actinotum*, 126; **anamorphum**, 126; *auberianoides*, 126; *clandestina*, 127; *terebrata*, 127  
*Odontia fimbriata*, 330  
*Oidium*, 223  
*Omphalia*, 165; *Austini*, 165; *Campanella*, 168, 250  
*Omphalopsis Campanella*, 168  
*Onagra*, 15; *biennis*, 15, 16, 30, 54, 64; *pallida*, 53  
*Onoclea*, 190; *sensibilis*, 190, 191  
*Onosmodium hispidissimum*, 11; *occidentale*, 10  
*Orchis spectabilis*, 9  
*Oropogon*, 152, 153; *loxensis*, 152-154  
*Oropogon loxensis* and its North American distribution, 152  
 Orton, C. R., Correlation between certain species of *Puccinia* and *Uromyces*, 194  
*Otomycosis aspergillina*, 279  
*Oxygraphis Cymbalaria*, 54, 64  
*Oxytropis*, 23; *Lamberti*, 23  
 Pacific Coast, Polyporaceae and Boletaceae of the, 91  
 Pacific Coast, The Agaricaceae of the—I, 205; II, 231; III, 294  
*Panicum virgatum*, 12, 202, 204  
*Pannaria*, 134; *leucosticta*, 134; *mollybdea*, 135; *virginosa*, 134  
*Panus stypticus*, 77  
 Parasite of catalpa, *Polystictus versicolor* as a wound, 263  
*Parmelia cetrata ciliosa*, 138; *cetrata subsidirosa*, 138; *laevigata*, 138; *perforata*, 138  
*Parmeliella tryptophylla*, 134  
*Paxillus*, 72, 243, 244; *atrotomentosus*, 244; *involutus*, 244; *ligneus*, 73; *pannoides*, 73  
*Peltigera canina*, 137; *polydactyla*, 137  
*Pentstemon alpinus*, 17, 31; *barbatus*, 17; *hirsutus*, 17, 31  
*Peridermium*, 50, 141-146, 184, 189, 190, 195, 282, 284; *abietinum*, 178; *balsameum*, 189-191; *carneum*, 29, 57; *Cerebrum*, 26, 143, 144; *coloradense*, 145; *Conorum-Piceae*, 146, 183, 184, 193; *consimile*, 178; *decolorans*, 178; *delicatulum*, 282, 284; *elatinum*, 145; *filamentosum*, 141-144; *fructigenum*, 140, 185, 186; *Harknesii*, 143, 144; **inconspicuum**, 283; *montanum*, 144, 284; *Peckii*, 186, 193; *pseudobalsameum*, 146; *stalactiforme*, 142; *Strobi*, 42, 143  
*Pertusaria cryptocarpa*, 137; *leioplaella*, 137; *tuberculifera*, 137; *velata*, 137  
*Pestalozzia*, 172, 331; *monochaeta*, 172  
*Petalostemon purpureus*, 10-12  
*Phacelia heterophylla*, 10  
*Phaeolus sistotremonoides*, 93  
*Phalaris arundinacea*, 11  
*Phialocybe*, 72  
*Philadelphus*, 63, 64; *coronarius*, 62, 63, 65  
*Phlebia*, 317; *hydroides*, 278  
*Phlox*, 29; *divaricata*, 12, 29  
*Pholiota*, 72, 244, 258; *aeruginosa*, 257; **albivelata**, 260; *anomala*, 259; *autumnalis*, 330; *candicans*, 259; *marginata*, 258; **McMurphyi**, 260; **oregonensis**, 262; *praecox*, 259; **subnigra**, 258; *unicolor*, 258; *ventricosa*, 260; **washingtonensis**, 259  
*Pholiotina*, 72  
*Phoma*, 106; *piciena*, 151; *pigmentivora*, 38  
*Phragmidium*, 194, 195, 197  
*Phragmites communis*, 54, 64  
*Phragmopyxis*, 282  
*Phryma leptostachya*, 9  
*Phyllospora furfuracea*, 131; *parvifolia*, 132  
*Physalis*, 11  
*Physcia*, 126; *comosa*, 140; *hypoleuca*, 140; *leucomela*, 140  
*Physcidia Wrightii*, 138  
*Physma byrsaeum*, 133  
*Picea*, 26, 178, 183, 184; *canadensis*, 177, 178, 183, 192, 193; *Engelmanni*, 145, 146; *excelsa*, 151; *Mariana*, 26, 33, 178, 183, 184, 192, 193; *Parryana*, 145; *rubra*, 151, 178, 192; *sitchensis*, 145

- Pilocarpon tricholoma*, 126
- Pinus*, 42; *chihuahuana*, 146; *con-torta*, 142-144; *Jeffreyi*, 143; *ponderosa*, 141, 143; *radiata*, 143; *rigida*, 282; *sabiniana*, 143; *taeda*, 29, 33, 57, 65; *virginiana*, 26, 33, 282, 284
- Pisum sativum*, 13
- Pleurage*, 116; *albicans*, 116; *lutea*, 116
- Pleuropus abortivus*, 6; *Earlei*, 332
- Pleurotus*, 4, 207, 217; *candidissimus*, 215; *corticatus*, 217; *densifolius*, 217; *dryinus*, 217; *geogenius*, 3; *oregonensis*, 217; *ostreatus*, 216; *sapidus*, 159, 216; *septicus*, 215; *serotinoideus*, 216; *serotinus*, 216; *subareolatus*, 217; *subsapidus*, 217; *subsepticus*, 217
- Plicaria echinospora*, 221
- Plowrightia*, 70
- Pluteolus*, 72, 74, 244, 246, 247; *californicus*, 246; *callistus*, 247; *expansus*, 247; *luteus*, 246, 247; *parvulus*, 246; *stramineus*, 246; *tropicalis*, 74
- Pluteus stercorarius*, 262
- Podiosoma*, 195
- Polemonium*, 29; *reptans*, 10, 12, 28, 29, 33
- Polygala Senega*, 12, 13
- Polygonatum commutatum*, 11
- Polyporaceae* and *Boletaceae* of the Pacific Coast, 91
- Polyporus carbonarius*, 217; *Caneus*, 109; *cutifractus*, 217; *elegans*, 93; *juniperinus*, 109, 110; *oregonensis*, 217; *perdelicatus*, 217; *Pseudotsugae*, 217; *sensibilis*, 217; *substipitatus*, 217
- Polystictus abietinus*, 113; *versicolor*, 263, 264, 266-270; *washingtonensis*, 217
- Polystictus versicolor* as a wound parasite of catalpa, 263
- Populus*, 30, 189; *acuminata*, 147; *angustifolia*, 147; *balsamifera*, 147; *grandidentata*, 147, 188, 193; *tremuloides*, 29, 33, 58, 65, 147; *trichocarpa*, 147
- Poria*, 113; *vaporaria*, 223
- Porodaedalea Pini*, 97, 105
- Preliminary notes on three rots of juniper, 109
- Prolifera* group, *Achlya* DeBaryana Humphrey and the, 319
- Psathyra*, 81
- Psathyrella*, 303
- Pseudotsuga*, 29, 30, 58, 261, 330; *mucronata*, 29, 33, 58, 65; *taxifolia*, 96
- Psilocybe*, 330
- Puccinellia airoides*, 54, 64
- Puccinia*, 9, 11, 14, 21, 22, 55, 56, 180, 181, 194-197, 199, 200, 203, 204; *Absinthii*, 21, 32; *albiperidia*, 13 (see page 49), 14, 15, 30 (see page 49), 49, 180, 193, 200, 201; *amphigena*, 18, 31; *Andropogonis*, 7, 17, 31; *angustata*, 7, 17, 31, 54, 64; *angustatoides*, 203; *Anthoxanthi*, 11; *argentata*, 20, 32; *Caricis*, 7, 17, 31; *Caricis-Asteris*, 16, 28, 30, 181, 193, 199, 204; *Caricis-erigerontis*, 199; *Caricis-Solidaginis*, 15, 30, 181, 193, 199; *cinerea*, 54, 64; *Crandallii*, 27, 33; *Distichlidis*, 11, 202, 204; *Eatoniae*, 61; *Eleocharidis*, 203; *Ellisiana*, 7, 9, 199, 200, 204; *emaculata*, 7; *fraxinata*, 7; *Gentianae*, 203; *graminis*, 39, 107; *Grossulariae*, 14; see page 49 for error; *heterantha*, 203; *Impatiensis*, 7; *Jamesiana*, 19, 31; *Lithospermi*, 28, 33; *Lygodesmiae*, 58, 65; *malvacearum*, 329; *monoica*, 61, 65; *montanensis*, 11; *Muhlenbergiae*, 18, 31; *opaca*, 203; *Opizii*, 16, 31; *Panici*, 202; *Pammellii*, 202-204; *Peckii*, 7, 15, 30, 54, 64, 181; *perminuta*, 9; *perplexans*, 179, 192; *Phragmites*, 54, 64; *poculiformis*, 7, 18, 31; *pustulata*, 7, 17, 31; *quadriflorula*, 28, 33; *Rhamni*, 11, 18, 31; *Ruelliae-Bourgai*, 203; *Seymouriana*, 19, 31; *Schedonardi*, 10, 53; *Stipae*, 19, 31, 59; *subnitens*, 18, 31, 52, 54-56, 64, 198, 204; *tosta*, 10, 52; *Triseti*, 59; *uniflorula*, 201, 204; *universalis*, 16, 31; *vexans*, 7; *virgata*, 10
- Puccinia* and *Uromyces*, Correlation between certain species of, 194
- Pucciniastrum agrimoniae*, 191; *arctium*, 190, 191; *minimum*, 184, 186, 193; *Myrtilli*, 186; *pustulatum*, 176, 177, 192
- Pulicaria*, 23
- Pyrola*, 183, 184; *americana*, 184, 193; *elliptica*, 184, 193
- Pyropolyporus Earlei*, 111; *Everhartii*, 104; *igniarius*, 97, 104; *juniperinus*, 110; *texanus*, 112
- Pyrus communis*, 25, 63
- Pyxine picta*, 139
- Quercus Ilex*, 172; *nigra*, 331; *Prinus*, 103; *rubra*, 26, 33, 170, 172, 331
- Ramalina denticulata*, 138; *linearis*, 138; *usneoides*, 138

- Ranunculus abortivus*, 61; *acris*, 179, 192  
*Razoumofskyia cryptopoda*, 142  
 Relatives, *Geaster minimus* Schw. and its. Notes on Iowa saprophytes—  
 I, 84  
*Resupinatus*, 207, 214; **atrocoeruleus**, 214  
*Rhamnus*, 11; *alnifolia*, 11, 18, 31; *cathartica*, 11  
*Rhododendron*, 150; *arborescens*, 150; *calendulaceum*, 150; *canadense*, 150; *canescens*, 150; *catawbiense*, 150; *nudiflorum*, 150; *maximum*, 150; *viscosum*, 150  
*Rhodora*, 184, 185; *canadense*, 184, 193  
*Rhynchospora*, 203  
*Ribes*, 14, 42, 180, 200, 201; *Cynosbati*, 9, 13, 30, 58, 180, 200; *floridum*, 28; *oxyacanthoides*, 180, 193; *prostratum*, 180, 193  
*Ricasolia subdissecta*, 136  
 Riddle, L. W., An enumeration of lichens collected by Clara Eaton Cummings in Jamaica—I, 125  
*Rinodina* *Conradi*, 139; *exigua*, 139; *sophodes exigua*, 139  
*Roestelia hyalina*, 50; *transformans*, 63  
*Rostkovites granulatus*, 100  
 Rots of juniper, Preliminary notes on three, 109  
*Rubus*, 190, 191; *allegheniensis*, 53  
*Rudbeckia laciniata*, 9, 12, 13, 28; *triloba*, 9-11  
*Ruellia*, 203  
*Rumex*, 196; *crispus*, 54, 64  
*Russula*, 207, 289; 291; *foetens*, 292; *emetica*, 290; *Mariae*, 290, 293; *obscura*, 291; **rubriochracea**, 293; *sericeomitens*, 290; **stricta**, 166; **sulcatipes**, 291; *uncialis*, 292  
 Rusts, Cultures of heteroecious, 175  
 Rusts, Two new species of, 282  
*Ryssospora*, 72  
*Salicornia*, 179; *europaea*, 179, 187, 192  
*Salix*, 147; *amygdaloides*, 147; *Bebbiana*, 147; *cordata lutea*, 147; *cordata Mackenziana*, 147; *discolor*, 188, 193; *fluvialis*, 147; *laevigata*, 147; *lasiandra*, 147; *lasiandra caudata*, 147; *lucida*, 147; *nigra*, 147; *Nuttallii*, 147; *rostrata*, 188; *sessifolia*, 147  
*Sambucus racemosa*, 180  
*Saprolegnia asterophora*, 326  
 Saprophytes, Notes on Iowa—I. *Geaster minimus* Schw. and its relatives, 84  
*Sarcobatus*, 198; *vermiculatus*, 54  
*Schedonnardus paniculatus*, 10, 53  
*Schizophyllum*, 213; *commune*, 213; *Egelingianum*, 214; *exiguum*, 214; *fasciatum*, 213; *flabellare*, 214; *mexicanum*, 214; *multifidum digitatum*, 214; *pavonium*, 214; *pusillum*, 214; *umbrinum*, 213, 214  
*Schizophyllum*, 213  
*Scirpus atrovirens*, 17, 31; *campestris paludosus*, 178, 192; *cyperinus*, 54, 64; *fluvialis*, 179; *paludosus*, 178; *validus*, 179  
*Scutiger oregonensis*, 93, 217; *reptipes*, 93  
 Seaver, F. J., The genus *Lamprospora*, with descriptions of two new species, 45; The genus *Lasiosphaeria*, 115  
*Senecio Douglasii*, 9, 11; *lugens*, 11, 20, 28, 31  
*Septoria ochroleuca*, 104, 170  
*Sidalcea oregana*, 10, 22, 53  
*Sitanion longifolium*, 18, 31  
*Smilax hispida*, 18, 31  
*Solidago*, 7, 16, 22, 181, 199; *bicolor*, 181, 192; *caesia*, 9; *canadensis*, 9, 13, 15, 19, 21, 31; *graminifolia*, 181, 192, 193; *lanceolata*, 15; *memoralis*, 15, 21; *rigida*, 10; *rugosa*, 15, 21, 32, 181, 193  
 Some western Uredineae which attack forest trees, Notes on, 141  
*Sorbus*, 25; *americana*, 25, 32, 57, 65; *aucuparia*, 57, 63  
*Spartina*, 11, 12, 186, 201; *cynosuroides*, 12, 29; *glabra alterniflora*, 186, 187, 193; *gracilis*, 201, 202, 204; *Michauxiana*, 12, 19, 29, 31, 33, 186, 187, 193, 201; *patens*, 12, 186, 187, 193  
 Spaulding, Perley, Notes upon tree diseases in the eastern states, 148  
 Species of *Puccinia* and *Uromyces*. Correlation between certain, 194  
 Species of rusts, Two new, 282  
 Species, The genus *Lamprospora*, with descriptions of two new, 45  
*Spergularia canadensis*, 179, 186, 187, 193  
*Sphaeria canescens*, 118; *emergens*, 119; *globularis*, 121; *hirsuta*, 119; *hispida*, 119; *mucida*, 118; *mutabilis*, 118; *orthotricha*, 119; *ovina*, 115, 117, 118; *Rhacodium*, 119, 121; *spermoides*, 121; *strigosa*, 118; *sublanosa*, 118; *terrestris*, 120  
*Sphaerophorus compressus*, 126  
*Sphaeropsis*, 34, 36; *tumefaciens*, 329  
*Spongipellis sensibilis*, 93, 217  
*Sporobolus asperifolius*, 10, 52; *neglectus*, 13

- Steccherinum, 309, 317; adustulum, 313; **basi-badium**, 314; **di-chroum**, 310, 311; **laeticolor**, 316, 318; Morgani, 315, 316; **multifidum**, 317, 318; **ochraceum**, 310-312, 317, 318; **Peckii**, 314, 315; **pusillum**, 313; **quercinum**, 277; **rawakense**, 312, 313, 315; Rhois, 310, 311, 314
- Steccherinum, The genus. Type studies in the Hydnaceae—II, 309
- Steironema, 12, 202; ciliatum, 12, 29, 201
- Stellaria graminea, 187; media, 187
- Stereocaulon cornutum, 133; pityrizans, 133; ramulosum, 133
- Stereum purpureum, 105
- Stevens, N. E., *Polystictus versicolor* as a wound parasite of catalpa, 263
- Sticta, 135; aurata, 136; crocata, 136; crocata leucosticta, 136; damacornis, 136; damacornis elongatolaciniata, 136; damacornis sinuosa, 136; dissecta, 135; dissecta corrosa, 135; erosa, 136; Fendleri, 135; Mougeotiana, 136; peltigera, 136; quercizans, 137; tomentosa, 137; Weigelii, 137
- Stipa, 19, 31, 59, 60; spartea, 19, 31; viridula, 59
- Strobilomyces strobilaceus, 99
- Stropharia, 294, 300, 305; aeruginosa, 300; bilamellata, 302; **longistriata**, 301; **magnivelaris**, 301; **semiglobata**, 2, 300, 301; **semigloboides**, 301; stercoraria, 300, 301
- Studies in the Hydnaceae, Type—I. The genus *Manina*, 271; II. The genus *Steccherinum*, 309
- Suedia, 179; maritima, 179
- Suillus luridus, 100, 163
- Symphoricarpos oreophilus, 28; pauciflorus, 10; racemosus, 10-12, 20, 27, 33
- Tachaphantium, 40
- Tapinia, 72, 243, 244; **lignea**, 73; panuoides, 244
- Teloschistes flavicans, 139
- Temperatures, *Aspergillus* infecting *Malacosoma* at high, 279
- Thalictrum dioicum, 9; polygamum, 10
- Thalloidima, 131
- Thelophora, 317; multifida, 317
- Thelotrema, 126; concretum, 127
- Thraustotheca, 87, 88; clavata, 87
- Thraustotheca clavata, 87
- Three rots of juniper, Preliminary notes on, 109
- Thuya, 95; plicata, 92
- Tissa canadensis, 11, 12, 19, 21, 55
- Tonina **janeirense**, 131
- Trametes Pini, 145; Sepium, 268
- Tree diseases in the eastern states, Notes upon, 148
- Trees, Notes on some western Uredineae which attack forest, 141
- Tremella, 40
- Tricella**, 282; **acuminata**, 282
- Tricholoma **jalapensis**, 332; **jamai-censis**, 332; **subisabellina**, 332
- Trifolium, 24; medium, 24, 56; pratense, 13, 24, 56; repens, 24, 56
- Trillium cernuum, 11
- Triphragmium, 195
- Trisetum, 59-61; majus, 11, 59, 65; subspicatum, 59, 60, 65
- Tropical agarics, New combinations for, 331
- Tropical North America, The Agaricaceae of—V, 72
- Tsuga, 184, 185, 189, 191; canadensis, 150, 176, 177, 182, 184-186, 188, 189, 191, 193; ciniastrium minimum, 185; heterophylla, 95; Pattoniana, 119
- Tubaria, 244, 249, 250; brevipes, 250; contraria, 249; deformata, 249; Eucalyptii, 249; furfuracea, 249; inquilina, 249; pallescens, 249; pellucida, 79; tenuis, 249
- Two new species of rusts, 282
- Two new species, The genus *Lamprospora*, with descriptions of, 45
- Type studies in the Hydnaceae—I. The genus *Manina*, 271; II. The genus *Steccherinum*, 309
- Tyromyces caesius, 94; **carbonarius**, 94, 217; **chioneus**, 94; **cutifractus**, 94, 217; **guttulatus**, 95; **perdelicatus**, 95, 217; **Pseudotsugae**, 95, 217; **sempileatus**, 96; **semisupinus**, 96; **substipitatus**, 96, 217
- Urecolaria, 127
- Uredineae in 1910, Cultures of, 7
- Uredineae in 1911, Cultures of, 49
- Uredineae which attack forest trees, Notes on some western, 141
- Uredinopsis, 190, 191; mirabilis, 190, 191; Pheopterididis, 191; Osmundae, 191
- Uredo, 194, 195, 197; Bigelowii, 147; Fabae, 194; Medusae, 147; Oxytropi, 23
- Uromyces, 12, 14, 21, 22, 55, 56, 179, 186, 194-197, 199, 201, 203, 204, 223; Astragali, 23, 32; acuminatus, 29, 33, 201, 202, 204; caricina, 22, 199; cucullatus, 203; Eleocharidis, 12, 203; graminicola, 12, 202, 204; Junci, 22, 32; Medicaginis, 24, 32,

- 56, 64; *Peckianus*, 12, 55, 56, 64, 179, 192, 198, 204; *pedatatus*, 200, 204; *perigynius*, 21, 22, 32, 181, 192, 199, 204; *plumbarius*, 203; *Rhynchosporae*, 203; *Ruelliae*, 203; *Scirpi*, 178, 179, 192; *Spartinae*, 12, 186, 187, 193; *speciosus*, 203; *Sporoboli*, 13; *Solidagini-Caricis*, 22; *uniporulus*, 14, 201, 204
- Uromyces*, Correlation between certain species of *Puccinia* and, 194
- Urtica gracilis*, 17, 28, 31
- Usnea angulata*, 138; *ceratina*, 138; *dasyopoga*, 139; *florida*, 139; *laevis*, 139; *plicata*, 139
- Ustilago esculenta*, 104
- Uvularia grandiflora*, 10, 11
- Vaccinium*, 146; *canadense*, 186; *pennsylvanicum*, 177, 186, 192
- Vaginata*, 231, 239; *farinosa*, 3; *vaginata*, 239; ***velosa***, 239
- Vagnera stellata*, 11
- Venenarius*, 231, 239; ***bivolvatus***, 241; ***calyptratoides***, 241; ***calyptratus***, 241, 242; ***mexicanus***, 332; *muscarius*, 240, 243; ***ocreatus***, 240; ***pantherinoides***, 242, 262; ***phaloides***, 240; ***praegemmatus***, 243, 262; ***solitarius***, 240; ***umbrinidiscus***, 242, 262
- Vernonia*, 29; *crinita*, 29, 33; *gigantea*, 57, 65
- Verticillium*, 104
- Viburnum*, 318
- Viola*, 199, 200; *septentrionalis*, 11
- Viorna Douglasii*, 11
- Vitis*, 34
- Volvaria* ***Bakeri***, 332; ***cubensis***, 332; ***Earlei***, 332; ***jamaicensis***, 332
- Volvariopsis* ***Bakeri***, 332; *cubensis*, 332; ***Earlei***, 332; *jamaicensis*, 332
- Western Uredineae which attack forest trees, Notes on some, 141
- Wound parasite of catalpa, *Polystictus versicolor* as a, 263
- Xanthoxylum americanum*, 10, 52, 53
- Zexmenia*, 203
- Zizania latifolia*, 104